
Basic Immunology Lecture 23rd

Immunological tolerance Cellular and molecular mechanisms of the immunological tolerance Autoimmunity



TOLERANCE - CENTRAL - PERIPHERAL: PASSIVE and ACTIVE

AUTOIMMUNITY - PHYSIOLOGIC REGULATION - AUTOIMMUNE DISEASES

Types of tolerance

- <u>Central tolerance</u> (selection of autoreactive T and B celle in the Thymus and Bone marrow)
- Peripheral tolerance
 - Lack of co-stimulation
 - Failure to encounter self antigens
 - Receipt of death signal
 - Control by regulatory T cells





Figure 1-14 Immunobiology, 6/e. (© Garland Science 2005)



Central T-cell -Tolerance thymic selection


Positive Selektion:

→ MHC RESTRICTION


Negative Selection:

→ <u>TOLERANCE</u>

Affinity model of thymocyte selection



Natural regulatory T-cell (Treg)




In Thymus presented antigens

- Own thymus antigens of epithelial cells, DC, Macrophages
- General cell antigens
- Extracellular antigens
- Medullar epithelial cells express other organ specific antigens (gens) → "promiscuous Gene expression through AIRE (Autoimmune regulator transcription factor)
- Infection related antigens


Central B-cell tolerance in BM


- 1. Receptor-Editing
- 2. Deletion with Apoptosis
- 3. Rezeptormodulation: BcR-downregulation \rightarrow Anergy

B cell selection in **BM**



Rezeptor-Editing





Passive tolerance


Unresponsiveness: no MHC recognition or inhibited cellular differentiation.


- Tolerance induced by the nature of the antigen
- Tolerance induced by the body

Passive tolerance induced by the nature of the antigen


- chemical nature
- dose of the antigen
 - low dose tolerance (T cell mediated, long ranging)
 - high dose tolerance (B cell mediated, short ranging)
- mode of the administration

b High-dose tolerance





Failed co-stimulation results low dose tolerance



Tolerance induced by the body

sequestered antigens

no MHC recognition no antigen presentation no systemic response

- heredited or acquired immunodeficiency
- clonal anergies
- induced tolerance


ACTIVE TOLERANCE

"Immunological homunculus"

- Low affinity IgM natural autoantibodies produced by CD5+ B cells
- γ/δ T cells
- Innate-like function


INDUCED Treg Cells

Oral tolerance




M-Cells - Endocytosis Epithelial cells – Transcytosis DC Through Tight Junctions

Development of induced Treg



Genetically well conserved antigens recognized by natural (auto)antibodies

Heatshock proteins	hsp65, hsp70, hsp90, ubiquitin
Enzymes	aldolase, citockrom c, SOD, NAPDH, citrate synthase, topoisomarase I.
Cell membrane components	β2-microglobulin, spectrin, acetylcholin receptor
Cytoplasmic components	actin, myosin, tubulin, myoglobin, myelin basic protein
Nuclear components	DNS, histones
Plasma proteins	albumin, IgG, transferrin
Cytokines,	IL-1, TNF, IFN, insulin,
hormones	thyreoglobin



AUTOIMMUNITY

- Physiological autoimmunity: part of the normal immunological regulation
 Natural autoantibodies: low affinity IgM produced by B1 cells
- Pathological autoimmunity: diseases caused by self reacting immune responses with <u>permanent</u> <u>tissue/organ injury</u>

High affinity IgG autoantibodies produced by T dependent B2 cells


Natural and pathologic autoimmunity

Natural autoantibodies

- Polyreactivity
- Low affinity
- Usually IgM
- ng/ml conc.
- products of CD5+ B1
 cells
- Target antigens: HSP, DNS, ACh R, (conservative structures)

Pathological autoantibodies

- High affinity
- IgG, IgA, IgM type antibodies
- mg/ml serum conc.
- products of Mature B2 type
- lymphocytes
- target antigens:
 - Cell surface structures,
 - receptors, proteins from the
 - cytoplasm, nucleoproteins



Autoimmunity

© copyright 2000 gary caviness/BioGrafix for more go to biogeek.com

Autoimmune diseases affect 5-7% of the population !

Autoimmunity by the failure of self tolerance


- Abnormal selection of lymphocyte repertoire
- Polyclonal activation of anergic selfreactive lymphocytes
- Stimulation by foreign antigens that cross-react with self

Pathomechanism of autoimmunity

- Inflammation and tissue necrosis
- Cellular components: (T cells CD8 and Th1, NK, Mf, DC, Ne, Eo, Ba, Mc)
 - Humoral components: (Ig+complement, ADCC, cytokines, chemokines, tissue hormones and mediators)

Autoimmune steady state

Self reacting immune response with tissues damages



Copyright ©2001 Shlomi Mallachi//smart3d

Active tolerance and tissue repair

Pathomechanism of autoimmunity

<u>Multifactor mechanism</u>


(general catastrophe of bio-regulation caused by external and internal factors)

- Autoimmune "steady state" (failure of dynamic balance on self tolerance and autoimmunity)


- Role of infections (molecular mimicry or inefficient natural antibody network)

Pathomechanisms of autoimmune diseases

- Autoimmunity by the antigen
- Failed differentiation and selection of lymphocytes
- Genetic background



The predictive relevance of autoantibodies




Pathogens and human antigens	Peptid residues	Overlaping sequences
Human cytomegalovirus IE2 HLA-DR molecule	79 60	PDP <u>LGRPD</u> ED VTE <u>LGRPD</u> AE
Poliovirus VP2	70	STT <u>KESRGT</u> T
Acetylcholine receptor	176	TVI <u>KESRGT</u> K
Papilloma virus E2	76	SLH <u>LESLKD</u> S
Insulin receptor	66	VYG <u>LESLKD</u> L
<i>Klebsiella pneumoniae</i> nitrogenase enzym HLA-B27 molecule	186 70	SR <u>QTDRED</u> E KA <u>QTDRED</u> L
Adenovirus 12 E1B	384	<u>LRRGMFRPSQCN</u>
Alfa-gliadin	206	LGQGS <u>FRPSQ</u> QN
HIV p24	160	<u>GVETTTPS</u>
Human IgG	466	<u>GVETTTPS</u>
Measles virus P3	31	<u>EISDNLGQE</u>
Myelin basic protein	61	<u>EIS</u> FK <u>LGQE</u>


Thyroid

Graves disease Hashimoto thyroiditis	DR3 DR5	3.7	TSH receptor ↑ Thyroid microsoma peroxidase, thyroglobin ↓
Pancreas IDDM	DR4/ DR3 DQB 0302	20 <u>100</u>	Beta island cells ↓ GAD, HSP60, junB, insulin, pre/pro insulin
Neural system Sclerosis multiplex Myasthenia gravis	DR2 DR3	4.8 2.5	Brain medulla, MBP, PLP, MOG, MAG Peripheral neurons- striated musceles Acetylcholine receptor
Hearth: rheumatic fever	DR3, DR4		S. B-haemolythicus M/myosin
Blood: AHA, thrombocytopenia			Vvs gP Thrombocyte gP

SLE	DR3/ DR2	5.8	Kidney, serous layers ds/ssDNS, Sm-IC, SSA
Sjögren syndrome			Exocrine glands, salivary glands, liver, kidney, brain, thyroid gland, hearth, lung, gut
Rheumatoid arthritis (RA)	DR4 DR1	4.2	Joint connective tissue, collagen Type II, IgG RF
Spondyloarthritis	B27	<u>90</u>	Vertebrate
(SPA) Reiter disease	B27	<u>33</u>	Clamydia, Yersinia
Salmonella/Shigella arthritis	B27	<u>20.7</u>	




Hashimoto's thyroiditis





Exophthalmos (bulging eyes)




Graves' disease is a common cause of hyperthyroidism, an over-production of thyroid hormone, which causes enlargement of the thyroid and other symptoms such as exophthalmos, heat intolerance and anxiety Normal thyroid

Enlarged thyroid




Diffuse goiter


STIMULATING AUTO-ANTIBODIES (Graves' disease)



Myasthenia gravis




Raynaud's Syndrome








Progressive Systemic Sclerosis



The artery shows early organization with "onion skin" change caused by lamellation and mucoid change with swelling of the intimal layer, with corrugation of the glomerular basement membrane. (Jones' silver stain, magnification X200).



Fibrous organization of the intimal injury of arteries in a more chronic stage of progressive systemic sclerosis . (Periodic acid Schiff reaction, magnification X400).

Rheumatoid arthritis

